

Birds on Top of 4/6 83.

Lunda cristata

Caprimulgus columbianus
Chordeiles ~~*caprimulgus*~~
Chordeiles ~~*caprimulgus*~~

Chordeiles cristatus

Mergus serrator

Fulica americana

~~*Larus borealis*~~ - 5 eggs?

Larus argentatus

Larus argentatus

Larus argentatus 3 eggs.

Larus argentatus

Phalaropus lobatus

Phalaropus lobatus

Birds at other locations


Scaphiopus *linaria*

Scaphiopus *fastidiosa*

Scaphiopus *fastidiosa*

Agave *complanata*

Measure of water for Tokyo 0.9-1.0 m.
length of pipe 4.0 m.



Storoccephalus dubowski

(no 2184)
Oscha shot one (2/683 between Luvina
and Saravuma; it was in company with
two of the same kind.

Storoccephalus dubowski

1883 The first one at the village 8/87. - The
following day an other. - The first pair
seen, when they were said to be
seen (B. 1883) 20th Aug 17. -
The first pair seen, and the first pair seen, seen
day. The first pair seen, seen. 20th
and seen from B. 1883.

March 1st
Sun 7

Chas. L. Smith

Stenella melanura

Several - 3 to 5 - near shore 4/5 83 and
also the following day, when I got two -
1 1/2 saw two individuals. - 12/6 saw two
about 11 am off of pier

They seem to be breeding this year
with us in some large trees into the center
of "Mesa Verde" 5/5 83 about 40 miles S. of D.

Stenella kermadecensis

The first one 455 at 10/5 85 - to 10/10 85. This
specimen was seen about 4/5 - 7521. seen near
at Tynagh C. D. - ~~Howe~~

None have been seen ^(10/5 83) since my return
from Petropavlovsk 3/5 83 -

Rev. R. J. ...

Procyon flavus
Young very abundant in Michigan
ground was shot on May 22nd, 23rd, 24th
and 31st of May 1883. Another was shot at
Ladysmith 3 of June 84. - Observed two
in Canada. Toronto 5/6. - Two seen
and one shot 10/6 83. -

Chelidon Hepuensis Ridgw.

3/6 83 one seen, but has probably arrived long
ago, as the eggs obtained 3/11 83 probably
belong to that species. - Met fully fledged
young at Karabulij, Ceylon Is., 9/7 83, and got a nest
with two eggs at Glinka 13/7 83, with two seen
to breed twice. -

Besides the ^{common} chirping and twitter when it has
a short song tsip, tsip, tsip, and its call
note is a short ~~tsip~~ tsip tsip.

2 15/4 83 2 in Karabulij, Ceylon Is. -
the other in Glinka 13/7 83 -

Erythronema leucura?

As I in the morning. 83 got
the first greenman on board the steamer
Alexander when on the way from Petro-
paulski to Derling Island, and showed
it to Dr. Dybowski, he asked me, whether
this species ~~did~~ occurs on the island
or not. As I had never seen or got
it there, I answered, that it did
not occur, having no idea of - of
course - that a ♀ was shot on
the island in the very hour in which
this conversation took place.

Muscicapa
haliae

A large number must have arrived
yesterday or last night (8/683) as about
10-12 were observed and 6 shot. Yester-
day the wind was S and changed during
the night to E with variable force from
very light to fresh. - The following day I
saw at least 9 and shot 5. - The 10th
2 were shot, and several seen. - Tomorrow
in all places sheltered from the high northern
wind of today (11/683), I myself counted
more than 20, four were shot. - 12/6 several
seen. - For the last two days they have not
been observed here at Gavane, but today (15/6)
Kasim brought some from Severin. - 17/6 I shot
a whole one at 200 yds. - Two seen, one of which
shot at Ostchansk Reichtler among the "Dunes"
19/683. - Orde brought one shot between Severin
and Savanna 21/683, and seen several. -

Felix Muscula

Chelidon saturata.

Cocher states, that he saw two swallows, black above and reddish brown below at the North Dockery D.I. 19/6 83, but as they were too near the seals it was impossible to secure them. - 22/6 83 old Sur. dukowski saw one swallow, black above and below, at Ladiginsk. -

Empidonax hammondi

Rather numerous this ~~day~~ spring; besides the 5 ones prepared, 3 others were received about 1/6 83. -

Acrocephalus genei

Sp. at Karachuk 7/183 about three.
fully fledged young in one shot. It
stands as if the parents were for a
second time, as I have met several
pairs in last pursuit of their love affairs.
They are usually found only along the steep
rocks of the ~~the~~ coast, but sometimes also
in the interior. Thus I met a family 10/183
at a small creek in the interior of Copper
Island at Karachuk; the note of the young
is deeper and lower than that of the adults
and is almost thrushlike. —

Leptoclebra brunneicollis

2087 2 1/2

Arctostaphylos

Isobrya variabilis?

Shot one example ~~just below~~ in the narrow valley just behind where the water supply of the village is. It was terribly injured by the shot (2 grain powder and shot about no. 2), so that it represented only a lower, single hole. One of both mandibles broken and the skull completely broken to pieces etc. It is a luck, that the color is so conspicuous, because the skin is only so good - or so bad - as the circumstances would allow -

— "Isobrya variabilis" Tay. J. p. Am. 1881 p. 124
from ~~Isobrya~~ and ~~Hammonia~~. —
Isobrya variabilis K. & C.

Hymenocera aureolus

Isobrya variabilis

Electrophagus variatus

1882. 13/3.83. Still only single birds. Commenced
to sing. - 18th April. Still only single birds. -
28th. The first flock consisting of 5 individuals
at Ruff. - 1st May. 1st. - Upper Is. natives assert, that
they lay large eggs twice in summer. -

Calcarius lapponicus

1885. The first was seen on the 22nd of April; the first in shot the following day.

— 14th. A fellow started from the shore today. The bird was plumbeous, rufous and tawny above, and the wings of a much lighter plumage.

Calcarius

Florida

1894-95 and the time he spent in the
field.

Cephus americanus

Brachyotus sculptus

- 4/5/81 Saw one at the old graveyard, N. I.
8/6/81 Found flowers of this in the old graveyard

Brachyotus sculptus

Alcedo canadensis

One seen on the river, N. of the falls, on July 15.
One seen in the neighborhood of the falls, on July 15.
One seen on the river, N. of the falls, on July 15.

One met a family of 5 indiv. on the road
between Seven Rookery and Gavan B. I. 3/983.

He says that a pair was seen the whole summer
in the neighborhood of the rookery, and that
Grebishky has got nest-youngs from Kameney,
some years ago. —

Alouatta palliata

Alouatta palliata

Fale

at a "Hole" near
several young ones, near Gavan, Copper Id.

Thalassartia pelagica

Found in Lough Neagh off Carr. Hill
on 25 Feb. 1891. at 10 fms. depth. The
specimens from Lough Neagh are the
young.

On 25 Feb. 1891

Halimedes

Lepus

of 13 ind. shot 29/5 87, but had one or
two dark patches, as usual the knee of
new feathers yet. - They are much more
difficult to shoot now than before, and
while a month ago 30-40 were the
average ~~number~~ result of half a day's
shooting, it is now only 10, that is
after 10. - 29/5 shot 14, a flight of 915 ft.
the day before and shot, were seen, but fallen.

April - 4/5 83 found a nest close to the Chalky
bank, at Sandpoint Landing, "Hill", Kentucky, in
a bushy place. The nest a diameter of 140 mm
consisting of dry grass and a few feathers.
The eggs were small, pale -

Spunta da letitica

Charadrius fuscus

1887. The first one was shot the 15th of May.
June 2^d. I observed half a dozen at the
Tomb and shot one. -

St. George

Hyperolius glaucus

9883. Copper Bl. Observed at least half a dozen
on the about 1000' high ridge between Cirsium and
Hemlock.

Hyperolius glaucus

4/6 83. Topokof Is. Nest with 3 eggs, ♀ shot on the
nest (no 2123) The eggs were laying with the
points inward, and downward ~~so~~ in a
slight hollow, ~~located~~ between four Sesuvia
arborescens, ~~with the~~ dry particles of leaves
and bark of this plant and numerous seeds of
the same formed the nest; they were evidently
brought together by the bird. Situation about
40 feet from ~~the sea~~ and about 14 feet
above the level of the sea. -- * high water mark.
One of the eggs was clear, the second contained
an embryo with large eyes, the third one
a still more developed one, with wings, legs
etc

8/783. *Cyprip. fl. Canadensis*, also there above 1000'. Seems to take the highest mountains where the young are fledged. — The call-note is a loud doo-riit! —

121-780 at Coppe Island Glinke. I was brought a not yet fully feathered young diving. Allowed to run on the floor, it immediately commenced a very hot pursuit of the flies, which were caught with a remarkable precision and rapidity. It did not take any notice of the presence of man, but when the dog in the corner room, it suddenly dropped flat on the floor with withdrawn neck, making itself as low as possible and rested so absolutely immovable until the dog had turned its back to it, when it ran off to any concealed place to hid itself. Within ^{half} an hour it had, however, made the experience that the dog did not ~~care~~ ^{take any notice of} it, and consequently ^{it's apartment} paid as

little attention to the dog as to the man. (no 2242).

Gallinule

18/3³³ With me too, the 1st and 2nd are -
I remember and mention more numerous
the young of the dog, the young of the
- I remember the 1st and 2nd of the
only, a number with the 1st and 2nd of the
of the sound of the 1st and 2nd of the
now, that the sound is passed through the
throat, but that the sound of the 1st and 2nd of the
causes by the vibrating motion into which
the body is brought when within the cage,
not in the same way as the 1st and 2nd of the
vibration of the 1st and 2nd of the
sound of the 1st and 2nd of the
body is a vibrating motion. The vibration
of the 1st and 2nd of the body, it is easily seen.

Stenocentrus maculatus

Thryothorus

1891. Found at ...
the ...
...
...

1892. ...
...

Atadionus thymenickii

Chrysophanes flavus

Hydrogaster quadrata

Sceloporus sublineatus

Two shot at Gavanski Owl D-I. 17/6 83.
brother of the same 22/6 83.

Sceloporus sublineatus

Elidora alpina

Alouatta palliata

Alouatta palliata

Thunbergia
One shot 20/6 and one 21/6 83 at (Severus), Emilia,
Novski days, D. I. Was alone. —

Thunbergia *Thunbergia* *Thunbergia* *Thunbergia*
and many others

Osche met a flock on the plateau between
Garam and Severus D.I. 3/983, feeding in the berries
of *Eupatorium*.

Lincoln's Sparrowhawk

no band of white on neck but low, w. feathers
15/5/83, in gray plumage.

1 shot and prepared during my absence

One of two shot at Gavan N.Y. 1/6/83, in red
plumage.

Two shot at Sverine 18/6/83.

One at Glinka, Copper Gt mid 7/83 and another
at Pestchaniy, 30/7/83.

Totipotency

Stegocryptus glauca

1. 1st & 2nd 7/6 83.-

1. 1st & 2nd 7/6 83.-

Ateneoetes nigrinus

Ocho saw and shot the first one B. 9. 83 $\frac{25}{5}$,
during my absence in Oymyakon.

In Oymyakon I saw one on the bank of the
"4" + 83. - which was seen in the Oymyakon
"2" + 83. - which was seen in the Oymyakon

Settia hyperborea

Settia

Limnocinclus acuminatus.
Longtail 1

Lepus sylvaticus

During my absence in Pel. gaudet
observed by the first house at Pel. gaudet
May 24th 1883. To the 1st of June I found them
numerous on the tundra.

Lepus sylvaticus

Chow — ?

Several white geese with black tipped
wings are reported to have been seen on
the surrounding lakes, northern part of Sering St.
5-6 Sept 1883. Said to be very shy.

Tringa allardii

allardii *Tringa*

Franta longicauda
9-24. Males in 3 at Singapore Dec 11-2
Catching from the sea. —

Eranta nigricans

Strophomena

Lepus chrysola

A single pair was found between 1880
and 1885. A 2. 24. of year 1883. Probably
a few were found in the Southern part.
Fossilized skeletons were found, that they
lived. One ♂ ~~observed~~ obtained 15 834
and another observed 5/681 on the Fossilized Gypsum.

Thys. lucida

They are common in the Southern part of the
country. It is not uncommon to find
19583 near the 1st of June in 1884.
19582 found on the 1st of June 1884.
In the Southern part.

Ophela acuta

23/87, a few ones observed between
Thurston and Palomares.

2/4/88. Quite low a small flock at Reef
to day. 28/4 10/4 and 2 at Larkspur.

4/6/88 Nest with 7 eggs on Top of Is.

10/4/88 a nest with 5 fresh eggs from same Is.

Ophela acuta

Mareca penelope

Order Nov 4 - Bristol - 1st of 1st order
to Nov 30/83. - ~~See above~~ The birds
rather numerous this spring, and several
were still seen 7/6 83.

Cyphopterus niger

Station 1350

April 13 1885. The first, a single one, was
seen in a small flock of black-backed goats.
The clouds the evening was fine and a
with a light easterly wind. - Some 200 or 300
was seen of C. hutchinsii by and of them.

22-24 of and common in the mountain valley
and the area of the valley.

Yellow and red of a large area
The large at the valley valley area
The at the valley valley area of the valley.

Quercus agrifolia

Enicetta falcata

Lynx "Enas falcata" Pall. Russ. T. III p. 701 n. 34.

Quell. hpt. N. I. n. p. 521. n. 86

Enas falcata Georgi Russ. p. 167. - Pall.

Europ. II p. 259 n. 343 " ex Maudsl.

Suppl. ad Europ. Russ. p. 21.

Enicetta falcata
Lynx falcata
D. J. 1833.

Fuligula cristata

7/685 a ♂ and a ♀ were shot in
Kamensk, only the ♂ being fit for pres-
ervation. Ochet thinks that he shot a ♂
during my absence in Petropaulskii
mistaking it for F. marila.

Fulix marila

575-83 the first of the series D. O. - by the
first one shot, 20.

Fuligula cristata

Arthropia

Mr Osche describes a duck which
he met 5/6 83 in Hannover in compar-
ison with Fuligula marila, as being
exactly like the male of this species,
but having a chestnut head. —

Clangula glaucion

Mr Osche describes it as being
exactly like the male of this species,
but having a chestnut head. —

Clangula glaucion

Thalassidroma pacifica

11/24/83 a large flock at sea.

21/6/83 I saw a large flock^{♂+♀} in the
small Creek Nolsjaja Rakuselnik, Severn.

Thalassidroma Brandt (see Thalassidroma)

Mem. Acad. Imp. Sc. St. Peterb. Vols. Sc. nat.
tom. six. (1848) - p. 4 Des.
φ δ υ α ε & histrio & υ η τ τ α and.

Thalassidroma pacifica

Arctia hyperborea
No. 1 of 1881 - 1st change of plumage.

Arctia hyperborea

Eulimnaria strobilata

1883. 12/3 Large flocks of ♂♂ and ♀♀ at Fritch
1/4 83 numerous in at 55. -
20-5/4 83 numerous flocks between
Kearney & Tuley. 1/4 83. -
few flocks also at 1/4 83 - 2/4 83 -
4/4 83 a few. - 5/4 83 a few. -
6/4 83 half a flock at 1/4 83 - 10/4 83
flocks between Kearney & Tuley. - Disappeared
about of May 83. -

Eulimnaria strobilata

Smilax angustifolia

185 to 211 from the bottom of the ...

See also

Melanetta americana
~~velutosa~~
Hakiston havde den fra Kautschatka.

Melanetta americana

Melanetta fusca
St. J. 1854

Mergus serrator

Copper Pl. Petrichanij Resolika, Gauran, 5883 a. p.

Mergus merganser

15/883! 2 - 1000 "marchants" cons. 1000 to 1000
at Gauran 1000

Cyprus

Larus glaucescens
had eggs on the tundra 16/1-83. —
found a bird was caught from
the same place. Photographed the bird
and eggs.

Arctic Skuas
April 1st 1880 - the first flock seen at
Tulavik.
First Young out 4883, Cape Mel.

Cissia brevicauda
April 12 1883 - La Jolla, Cal. - common at
Toluca Lake

Cissia brevicauda

Sterna paradisica Mun. 22
the first ones seen and two shot
5/6 83 at the tundra at Gavan. 9/6 Osler saw
4 there.

Sterna longipennis
Besides the one shot several
others were seen. 21/6 83 one got from Emilia,
near Ki Lys D. I.

29/6 83 plenty of them at Rakovaja
Ancheta, Ancheta Bay, Kamt.
Ch. Kittitz ± p 322 *Sterna glacialis*,
Kitt. = *S. kamoharui* Penn & Ball.

Calcearia lapponicus

Stenonema longicaudatus

4/6 83. One or two were observed
when on Tokoro Island. Got one
the next day. - Also one 2/6 83 from
the small lake at Marochin between
Sarama and Gavan B.I. - One from
Gladkovskij, Coppe Is. 12 83, and
then saw another at Karabelnij some days after --
At my arrival at Glinka. I found there two
specimens: tarsus blue, feet black. - In no 13
at Gavan Coppe Is. 12 83.

Stenonema parvicaudatus

4/5 83 The first one at Bay B.I.
7/6 83 An example of the light phase. 20/6 83.

Pinnipedia

1982. Found the remains of an adult seal
skull on the shore of the bay.
this year.

Dr. Clark saw one at 11/13/82, and not
less than 8 white ones 11/4/84.
I saw one adult 2/5/84 at Key 12-2.

Oceanodroma furcata

Metopidroma nelsoni

Fulmaridae

April 15 1982. In several places to the bay from 1981.
They appeared to be the same as the ones from 1981.

Phalaropus lobatus

Tringa borealis

Thalassidroma leucorhoa

Canterbury, England, 1883. ~~18~~ 1883. The
the same, but says, also that some in 1883.
Gann 1883. 1883. 1883. 1883. 1883. —

16 83. 1883. 1883. 1883. 1883. 1883. —
only one being in the nest. —

— 20/7 83. 1883. 1883. 1883. 1883. 1883. —
one pair at Portcharter Inlet, Gann.

Salicaria concolorata

Thymus aureus

Leucis viridula

7/5/81. Several returned from an expedition to
the River, during which he had observed
Spots, between the two ferns, I. 2.
The same, I think, is probably the same
the bird is smaller, more brown and blue,
not so abundant, and a little more varied
with brown, except the tail, which is light
and, as a rule, more curved than in the adult.
The nest was made of moss and some other
- 18/5/81. The first egg - and only one - was
seen, found in the nest. - 15/6. In 2 eggs
the first, - 27/8. and one more, I think
the first, the second, and the third, I think
they were eggs of the same species.

Leucis viridula

Limonium pycnanthum
7541 got one from ~~Long Island~~, *Peru*, *Peru*

Chalcid pycnanthum
38/4 88 - 3 at Long Island, Off. I.
4509 early in the morning plenty of *Chalcid* *pycnanthum*
- *Chalcid* *pycnanthum*, that if *Chalcid* occurs at
all in *pycnanthum*, it will surely be very
small, and *Chalcid* *pycnanthum* than *Chalcid* *pycnanthum*,
and *Chalcid* *pycnanthum* *Chalcid* *pycnanthum* *Chalcid* *pycnanthum*
there is a *Chalcid* *pycnanthum* *Chalcid* *pycnanthum* *Chalcid* *pycnanthum*
say that *Chalcid* *pycnanthum* *Chalcid* *pycnanthum* *Chalcid* *pycnanthum*

Chalcid

Syntherisma argentea.
1881. Several hundred small dark ones seen at sea
today, Oct. -

Cyprina edulis
Mr. Wm. Quinn, the fish dealer
of New York City, 1758 - 1853. several
at Topsham 4/683 in smaller or larger flocks.

Stomatopoda

Uria

As this will have been seen on May 1st
at Long Island Sound N.Y. I want to put down
some things at this time for when we cross to day
— When sitting on the rock shelves, the
Uria ^{usually} turn their breast to the rock wall and then
back to the sea, while Larus cirrhatus and
Fratriscula corniculata front the sea. ~~usually~~ Their
position of the latter is surely upright although
not so straight as that of the Uria
— Uria Uria

Uria Uria Uria

Chrysomelidae

Thyridopsis borealis

1882

